

Measurement of event shapes and α_s in e^+e^-

C. Pahl

*Max-Planck-Institut für Physik, Föhringer Ring 6, D-80805 Munich, Germany
Excellence Cluster Universe, TU München, Boltzmannstr. 2, D-85748 Garching, Germany.*

We report results on measurements of the strong coupling α_s from event shape data at PETRA and LEP. These include analyses of their distributions employing the recent NNLO calculations and analyses of their moments using NLO and power correction predictions.

1. INTRODUCTION

In an hadronic event the annihilation of the e^+e^- pair into an Z^0/γ and its subsequent decay into an quark pair is followed by gluon radiation and other processes. At an energy scale of about 1 GeV hadronisation takes place and the partons transform into hadrons. The process of hadronisation can only be described by Monte Carlo models or – more recently – analytically by power correction models. The ultimate goal is to measure the strong coupling; i.e. to test its energy evolution as described by QCD and to extract a value at some definite energy scale, commonly taken as the mass of the Z^0 boson. The most recent world average is $\alpha_s(M_{Z^0}) = 0.1198 \pm 0.0010$ [1].

The construction of the e^+e^- collider experiments JADE and OPAL is quite similar and a consistent measurement can be expected by using both. OPAL operated at centre-of-mass energies from 91 to 209 GeV, JADE at lower energy scales from 12 to 44 GeV, where the strong coupling and its energy evolution are stronger. The experiment ALEPH was one of the four LEP experiments.

The studied event shapes are thrust T or $1 - T$, C-parameter C and total jet broadening B_T ; these variables are sensitive to the whole event. Further heavy jet mass M_H , wide jet broadening B_W and Durham two-jet flip parameter y_{23}^D ; these are sensitive to only one suitable chosen hemisphere of the event. For the definitions of the variables we refer to [2].

2. NNLO ANALYSES OF EVENT SHAPE DISTRIBUTIONS

The recently completed NNLO calculations [3] have been fitted to event shape distributions measured by JADE [4] and ALEPH [5]. Figure 1 shows the fits in case of thrust; JADE measured $1 - T$ at centre-of-mass energies $\sqrt{s} = 14$ to 44 GeV, while ALEPH measured T at 91 to 206 GeV. In the JADE analysis resummed next-to-next-to-leading logarithms (NLLA) are included employing the $\ln R$ -matching scheme. In contrast to NLO analyses, these predictions fit well over virtually the whole phase space.

The major uncertainty of NLO analyses [2, 6] comes from varying the renormalisation scale μ_R for estimating the contribution of missing higher order perturbative terms. Therefore the dependency of the fit results on the scale factor $x_\mu \equiv \mu_R/\sqrt{s}$ has been studied. To compare with the NNLO+NLLA analysis, NNLO, NLO and NLO+NLLA studies have been pursued by JADE. ALEPH compares the NNLO analysis with NLO and NLO+NLLA. The analyses including NNLO show the smallest dependency on x_μ , which leads to a small scale uncertainty. The resulting $\alpha_s(M_{Z^0})$ values – combined from the variables thrust, M_H , B_T , B_W , C , y_{23}^D at all analysed energy points – are shown in figure 2. The values and their scatter from the different variables get smaller by the inclusion of the NNLO terms. The same holds for the inclusion of resummed logarithms. The JADE value from the NNLO+NLLA analysis is identical to the value from NLO+NLLA, but scatter and error are reduced. The theoretical scale uncertainties are considerably reduced by the NNLO terms. The ALEPH results are similar to the JADE results with somewhat higher central values. Combining the $\alpha_s(M_{Z^0})$ values from the NNLO analyses of the different variables, the energy evolution agrees with the QCD prediction of asymptotic freedom. The evolution is stronger in the JADE energy range, but the errors are smaller from the ALEPH points. The hadronisation effects scale with inverse powers of




Figure 1: Fits of the NNLO+NLLA prediction to JADE data (prelim.) and fits of the NNLO prediction to ALEPH data. The symbols show the measurement on hadron level with statistical errors. The fit ranges are indicated by arrows resp. solid lines.




Figure 2: $\alpha_s(M_{Z^0})$ results from the JADE or ALEPH analyses of $1 - T$ or T , M_H , B_T , B_W , C , $y_{23}^D \equiv y_3$, and combination of these variables. The results from the different energy points have been combined.

$\sqrt{s}$, so they – and their errors – are smaller at high energies. Therefore the ALEPH result, $\alpha_s(M_{Z^0}) = 0.1240 \pm 0.0008(\text{stat.}) \pm 0.0010(\text{exp.}) \pm 0.0011(\text{had.}) \pm 0.0029(\text{theo.}) = 0.1240 \pm 0.0033$ is more precise than the JADE result, $\alpha_s(M_{Z^0}) = 0.1172 \pm 0.0006(\text{stat.}) \pm 0.0020(\text{exp.}) \pm 0.0035(\text{had.}) \pm 0.0030(\text{theo.}) = 0.1172 \pm 0.0051$.

3. NLO ANALYSES OF EVENT SHAPE MOMENTS

The n -th moment of the distribution of event shape variable y is defined as

$$\langle y^n \rangle = \frac{1}{\sigma_{\text{tot.}}} \int y^n \frac{d\sigma}{dy} dy,$$

The moments have been measured for order $n = 1$ to 5 from JADE [7] and OPAL [2] data. With increasing moment order the peak of the integrand is shifted towards the multi jet region, so higher moments are sensitive to this region.

Employing hadronisation corrections from Monte Carlo models the data are compared [2, 7] with the NLO prediction $\langle y^n \rangle = \mathcal{A}_n \alpha_s(s) + \mathcal{B}_n \alpha_s^2(s)$ (schematically; terms accounting for correct normalisation and scale dependency are suppressed). The $\langle (1 - T)^n \rangle$, $\langle C^n \rangle$ and $\langle B_T^n \rangle$ predictions fit the data well. In case of the one-hemisphere observables $\langle B_W^n \rangle$, $\langle (y_{23}^D)^n \rangle$ and $\langle M_H^n \rangle$, the agreement is not good at low energies (but still $\chi^2/\text{d.o.f.} \simeq 10$). This effect can be understood from corresponding analyses of distributions, see figure 3. The $\mathcal{O}(\alpha_s^2)$ coefficient is negative over a large




Figure 3: Distributions of the one-hemisphere variables M_H , B_W and $y_{23} \equiv y_{23}^D$ at 14 GeV; JADE measurement and various fit curves [6]. As indicated, the dash-dotted curve is to be compared with our analysis. The range where this prediction is negative (but not plotted) is marked with an oval starting at zero on the abscissa.

part of the two jet range. So at low energies – where it contributes strongly – the distribution is unphysically negative. As the moment coefficient is the integral over the distribution coefficients, it can be judged as “unphysically low” – i.e. large corrections of higher order can be expected.¹ Therefore the NLO prediction of $\langle M_H^1 \rangle$ can not be fitted [2] and the description is very bad for $\langle B_W^1 \rangle$ – so this fit is not further used.

The fit results are increasing significantly and systematically with moment order for the two-hemisphere variables. The K-factors $\mathcal{B}_n/\mathcal{A}_n$ of the predictions qualitatively show the same behaviour. So the trend of large higher order contributions for the two-hemisphere moments seems to continue; i.e. in the NLO prediction a large part is missing, and this is compensated by an increased $\alpha_s(M_{Z^0})$ value. Combining the results from fitting the more complete predictions (i.e. those satisfying $\mathcal{B}_n < 0.5 \cdot \mathcal{A}_n$) gives $\alpha_s(M_{Z^0}) = 0.1262 \pm 0.0006(\text{stat.}) \pm 0.0010(\text{exp.}) \pm 0.0007(\text{had.}) \pm 0.0064(\text{theo.})$, consistent with the world average.

Hadronisation can also be described by analytical, *non perturbative* models. We tested [9] several of them with event shape moments data. The *dispersive model* by Dokshitzer et al. predicts a simple shift of the perturbatively calculated distribution when going from parton to hadron level, $d\sigma/dy = d\sigma_{\text{part.}}/dy(y - a_y \cdot \mathcal{P}(\alpha_0))$. Here the power correction factor a_y depends on the observable, while $\mathcal{P}(\alpha_0)$ does not. This description can be integrated giving predictions for the moments. The data fit these predictions well. The high $\alpha_s(M_{Z^0})$ values from the two-hemisphere

¹In fact the negative region is smaller in the NNLO prediction [8].

moments are not cured. The parameter α_0 should be universal – this is not the case especially for the values from $\langle B_W^n \rangle$, where this problem has been known from a study of the distribution [10]. Combining the values from fitting the more complete descriptions gives $\alpha_s(M_{Z^0}) = 0.1174 \pm 0.0050(\text{tot.})$, $\alpha_0(\mu_I) = 0.484 \pm 0.053(\text{tot.})$.

4. SUMMARY AND OUTLOOK

The NNLO and NNLO+NLLA fits of event shape distributions measured by JADE and ALEPH show significantly reduced scale uncertainty and reduced scatter from the different variables. The predictions fit well over a larger range of the distributions. The strong coupling was measured with a precision of 3% by ALEPH, $\alpha_s(M_{Z^0}) = 0.1240 \pm 0.0033$. Moments of event shape distributions have been measured by JADE and OPAL. The perturbative NLO prediction is adequate only for some moments; this incomplete perturbative description shows up in all studied non perturbative models. When passing from first to higher moments, new perturbative and non perturbative problems appear.

Better resummation is available: NNNLA has been calculated for thrust in soft collinear effective field theory [11]. The NNLO predictions for the moments are almost complete [12].

Discussion: Giulia Zanderighi – In a recent paper, Weinzierl calculated somewhat different NNLO coefficients than Gehrmann et al. Employing these calculations the α_s values are not expected to change much; but the scatter from the different variables could become still smaller. Chris Maxwell – An analysis of event shape moments within renormalisation group improved perturbation theory would be interesting. This way, the DELPHI collaboration was able to describe mean values of event shapes without any power corrections.

References

- [1] S. Bethke. Experimental tests of asymptotic freedom. *Prog. Part. Nucl. Phys.*, 58:351, 2006.
- [2] G. Abbiendi et al. Measurement of event shape distributions and moments in $e^+ e^- \rightarrow \text{hadrons}$ at 91 GeV - 209 GeV and a determination of α_s . *Eur. Phys. J.*, C40:287, 2005.
- [3] A. Gehrmann-De Ridder, T. Gehrmann, E. W. N. Glover, and G. Heinrich. NNLO corrections to event shapes in e^+e^- annihilation. *JHEP*, 12:094, 2007.
- [4] S. Bethke, St. Kluth, C. Pahl, J. Schieck. Determination of the strong coupling α_s from hadronic event shapes and NNLO QCD predictions using JADE data. *Submitted to Eur. Phys. J.*, C, 2008.
- [5] G. Dissertori et al. *JHEP*, 0802:040, 2008.
- [6] P. Movilla Fernández. Studien zur Quantenchromodynamik und Messung der starken Kopplungskonstanten α_s bei $\sqrt{s}=14\text{--}44\text{ GeV}$ mit dem JADE-Detektor. *PhD thesis*, 2003. RWTH Aachen.
- [7] C. Pahl, S. Bethke, St. Kluth, J. Schieck. Study of moments of event shapes and a determination of α_s using e^+e^- annihilation data from JADE. *Submitted to Eur. Phys. J.*, C, 2008.
- [8] T. Gehrmann. Status of NNLO jet calculations. In *Proceedings of FRIF workshop on first principles non-perturbative QCD of hadron jets, LP THE, Paris, France*, 12-14 Jan 2006.
- [9] C. Pahl. Untersuchung perturbativer und nichtperturbativer Struktur der Momente hadronischer Ereignisformvariablen mit den Experimenten JADE und OPAL. *PhD thesis*, <http://nbn-resolving.de/urn:nbn:de:bvb:91-diss-20070906-627360-1-2>, 2007. TU München.
- [10] P. Movilla Fernández. Event shapes from JADE data and studies of power corrections. *Nucl. Phys. Proc. Suppl.*, 74:384, 1999.
- [11] T. Becher, M. D. Schwartz. A precise determination of α_s from LEP thrust data using effective field theory. *JHEP*, 07:034, 2008.
- [12] T. Gehrmann. *Private communication*.